arXiv:1302.3051v1 [math.NT] 13 Feb 2013

Some Properties of Generalized Self-reciprocal Polynomials over Finite Fields

Ryul Kim [1], Ok Hyon SONG [1], Hyon Chol Li[2]

[1] *Department of Mathematics*
Kim Il Sung University, Pyongyang, D.P.R.Korea
[2] *Department of Mathematics*
Chongjin Education University No 2, Chongjin, D.P.R.Korea

January 22, 2013

Abstract

Numerous results on self-reciprocal polynomials over finite fields have been studied. In this paper we generalize some of these to a-self reciprocal polynomials defined in [4]. We consider the properties for the divisibility of a-reciprocal polynomials, estimate the number of all nontrivial a-self reciprocal irreducible monic polynomials and characterize the parity of the number of irreducible factors for a-self reciprocal polynomials over finite fields of odd characteristic.

Keywords: Finite fields, Self-reciprocal polynomials

1 Introduction

Self-reciprocal polynomials over finite fields are of interest both from a theoretical and a practical viewpoint, so they are widely studied by many authors. Let $S_q(n)$ denote the number of self-reciprocal irreducible monic(srim) polynomials of degree $2n$ over $\mathbf{F}_q$. Carlitz proposed the following formula and Meyn gave a simpler proof of it; see [3, 6, 8].

$$S_q(n) = \begin{cases} \frac{1}{2n}\left(q^n - 1\right), & \text{if } q \text{ is odd and } n = 2^s, \\ \frac{1}{2n}\sum_{\substack{d|n \\ d \text{ odd}}} \mu(d) q^{n/d}, & \text{otherwise} \end{cases} \tag{1}$$

Using the Stickelberger-Swan theorem, Ahmadi and Vega characterized the parity of the number of irreducible factors of a self-reciprocal polynomials over finite fields; see [2]. In [10], Yucas and Mullen classified srim polynomials based on their orders and considered the weight of srim polynomials. The problems concerning to the existence of srim pentanomials and the number of srim polynomials with prescribed coefficients have also been considered, see [1, 5]. On the other hand, in [4] Fitzgerald and Yucas introduced and characterized the generalized reciprocal polynomials to gave new descriptions of the factors of Dickson polynomials over finite fields.

In this work we find some properties for the divisibility of generalized reciprocal and generalized self-reciprocal polynomials. We also study the number of generalized srim polynomials and the parity of the number of irreducible factors of generalized self-reciprocal polynomials.

Throughout, $q = p^e$ be an odd prime power, $\mathbf{F}_q$ be a finite field containing q elements and $0 \neq a \in \mathbf{F}_q$. For $f(x) \in \mathbf{F}_q$, a monic polynomial of degree n with $f(0) \neq 0$, $\hat{f}_a(x) := \frac{x^n}{f(0)} f\left(\frac{a}{x}\right)$ is called *a-reciprocal* of $f(x)$[4]. That is, if $f(x) = \sum_{i=0}^n b_i x^i$ then $\hat{f}_a(x) = (1/b_0) \sum_{i=0}^n b_i a^i x^{n-i}$. $\hat{f}_a(x)$ is also monic and if α is a root of $f(x)$ then a/α is a root of $\hat{f}_a(x)$. The case $a = 1$ is a usual reciprocal in [2, 8, 10].

2 Divisibility of generalized reciprocal polynomials

In this section we summarize some properties for the divisibility of generalized reciprocal and generalized self-reciprocal polynomials over $\mathbf{F}_q$.

Theorem 1 *a-reciprocal of a product of two polynomials is the product of a-reciprocals of given polynomials.*

Proof. Let $f(x) = \sum_{i=0}^n b_i x^i, g(x) = \sum_{j=0}^m c_j x^j (b_0 \neq 0, c \neq 0)$. From the definition,

$$\hat{f}_a(x) = \frac{1}{b_0} \sum_{i=0}^n b_i a^i x^{n-i} = \frac{1}{b_0} \sum_{i=0}^n b_{n-i} a^{n-i} x^i,$$

$$\hat{g}_a(x) = \frac{1}{c_0} \sum_{j=0}^m c_j a^j x^{m-j} = \frac{1}{c_0} \sum_{j=0}^m c_{m-j} a^{m-j} x^j,$$

$$\begin{aligned} \hat{f}_a(x) \cdot \hat{g}_a(x) &= \frac{1}{b_0 c_0} \sum_{k=0}^{n+m} \left[\sum_{i+j=k} \left(b_{n-i} a^{n-i}\right) \left(c_{m-j} a^{m-j}\right) \right] x^k \\ &= \frac{1}{b_0 c_0} \sum_{k=0}^{n+m} \left(\sum_{i+j=k} b_{n-i} \cdot c_{m-j} \cdot a^{n+m-k} \right) x^k \end{aligned}$$

and

$$f(x) \cdot g(x) = \sum_{k=0}^{n+m} \left(\sum_{i+j=k} b_i c_j \right) x^k = \sum_{k=0}^{n+m} d_k x^k \left(d_k := \sum_{i+j=k} b_i c_j \right).$$

Therefore

$$\widehat{fg}_a(x) = \frac{1}{b_0c_0}\sum_{k=0}^{n+m} d_k a^k x^{n+m-k} = \frac{1}{b_0c_0}\sum_{k=0}^{n+m} d_{n+m-k} a^{n+m-k} x^k$$

We set $i' := n-i, j' := m-j$, then

$$d_{n+m-k} = \sum_{i'+j'=k} b_{n-i'} c_{m-j'}$$

and thus

$$\widehat{fg}_a(x) = \frac{1}{b_0c_0}\sum_{k=0}^{n+m}\left(\sum_{i'+j'=k} b_{n-i'}\cdot c_{m-j'}\cdot a^{n+m-k}\right)x^k. \qquad \Box$$

It might be $\hat{f}_a(x) = \hat{f}_b(x)$ for nonzero distinct elements a, b in $\mathbf{F}_q$. For example, if $f(x) = x^2 + c \in \mathbf{F}_5[x]$, $c \neq 0$, then $\hat{f}_2(x) = \hat{f}_3(x)$.

A monic polynomial $f(x) \in \mathbf{F}_q[x]$ is said to be *a-self reciprocal* when $\hat{f}_a(x) = f_a(x)$. In [4], the notion of a-self reciprocal polynomial has been defined for only even degree. Note that $f(x) = \sum_{i=0}^n b_i x^i$ is a-self reciprocal if and only if for each i, $b_{n-i}b_0 = b_i a^i$. When $i = n$, we see that $b_0^2 = a^n$.

First consider a case when n is odd. If a^n and thus a is not a square in $\mathbf{F}_q$, then there is not any a-self reciprocal monic(srm) polynomial of degree n. When $0 \neq a \in \mathbf{F}_q$ is a square in $\mathbf{F}_q$, if $f(x)$ is a-srm polynomial then either $b_0 = \sqrt{a}^n$ or $b_0 = -\sqrt{a}^n$.

Theorem 2 *Let n be an odd, $0 \neq a \in \mathbf{F}_q$ be a square in $\mathbf{F}_q$ and $f(x) \in \mathbf{F}_q$ be a-srm polynomial of degree n.*

(1) If $b_0 = \sqrt{a}^n$ then $f(x)$ is divided by $x + \sqrt{a}$.

(2) If $b_0 = -\sqrt{a}^n$ then $f(x)$ is divided by $x - \sqrt{a}$.

Proof. We have

$$\begin{aligned}
f\left(-\sqrt{a}\right) &= \left(-\sqrt{a}\right)^n + b_{n-1}\left(-\sqrt{a}\right)^{n-1} + \cdots + b_1\left(-\sqrt{a}\right) + \left(\sqrt{a}\right)^n \\
&= b_{n-1}\left(-\sqrt{a}\right)^{n-1} + \cdots + b_1\left(-\sqrt{a}\right), \\
\hat{f}_a\left(-\sqrt{a}\right) &= -\frac{1}{\left(-\sqrt{a}\right)^n}\left[\sqrt{a}^n\left(-\sqrt{a}\right)^n + \cdots + b_{n-1}a^{n-1}\left(-\sqrt{a}\right) + a^n\right] \\
&= -\left[\sqrt{a}^n + b_1\left(-\sqrt{a}\right) + \cdots + b_{n-1}\left(-\sqrt{a}\right)^{n-1} + \left(-\sqrt{a}\right)^n\right] \\
&= -\left[b_1\left(-\sqrt{a}\right) + \cdots + b_{n-1}\left(-\sqrt{a}\right)^{n-1}\right] = -f\left(-\sqrt{a}\right)
\end{aligned}$$

and from $f = \hat{f}_a$, $f\left(-\sqrt{a}\right) = -f\left(-\sqrt{a}\right)$ which implies that $f\left(-\sqrt{a}\right) = 0$. The second case follows similarly. $\Box$

Corollary 1 *If $0 \neq a \in \mathbf{F}_q$ is a square in $\mathbf{F}_q$, then $x+\sqrt{a}$ and $x-\sqrt{a}$ are only a-srim polynomials of odd degree over $\mathbf{F}_q$.*

Next let n be an even, that is, $n = 2m$. $f(x) = \sum_{i=0}^{2m} b_i x^i$ is said to be *trivial* or *nontrivial* respectively, according to $b_0 = -a^m$ or $b_0 = a^m$[4]. If $f(x)$ is a trivial a-srm polynomial, then $f\left(\sqrt{a}\right) = f\left(-\sqrt{a}\right) = 0$, hence $f(x)$ is a multiple of $x^2 - a$. Therefore $x^2 - a$ is only a trivial a-srim polynomial. From the definition, $b_m b_0 = b_m a^m$ and $b_0 = -a^m$, so that $b_m = 0$, hence trivial a-srm polynomials have always even terms.

Lemma 1 *$x^2 - a$ is a trivial a-srm polynomial and $\left(x^2-a\right)^2$ is a nontrivial one.*

Lemma 2 *The product of two a-srm polynomials satisfies the following multiplication table.*

$\cdot$	*trivial*	*nontrivial*
trivial	*nontrivial*	*trivial*
nontrivial	*trivial*	*nontrivial*

The proof of above lemmas are very simple so we omit.

Theorem 3 *If $f(x) \in \mathbf{F}_q[x]$ is a-srm polynomial of even degree, then it can be written as $f(x) = \left(x^2-a\right)^k \cdot g(x)$, where $g(x)$ is a nontrivial a-srm polynomial not divided by $x^2 - a$ and*

$$k = \begin{cases} \textit{odd}, & f(x) \textit{is trivial}, \\ \textit{even}, & f(x) \textit{is nontrivial} \end{cases}$$

Proof. Suppose that $x^2 - a$ divides $f(x)$. Then $f(x) = \left(x^2-a\right) \cdot f_1(x)$ for some $f_1(x) \in \mathbf{F}_q[x]$. If $f(x)$ is trivial, then $f_1(x)$ is nontrivial by Theorem 1 and above lemmas. And if $f(x)$ is nontrivial, then $f_1(x)$ is trivial, so it is divided by $x^2 - a$ and it can be written as $f(x) = \left(x^2-a\right)^2 \cdot f_2(x)$. Again from above lemmas $f_2(x)$ is nontrivial. Continuing this procedure implies the claim. □

If $f(x) \in \mathbf{F}_q[x]$ is divided by $x^2 - a$, then $f(x)$ has its roots $\sqrt{a}$ and $-\sqrt{a}$. If a is a square in $\mathbf{F}_q[x]$, then there might be a-srm polynomial which is not divided by $x^2 - a$ but has a root $\sqrt{a}$ or $-\sqrt{a}$. For example, if $q = 5, a = 4$, then $\sqrt{a} = 2, -\sqrt{a} = 3$,

$$f(x) = x^6 + x^5 + 3x^4 + 4x^3 + 2x^2 + x + 4 \in \mathbf{F}_5[x]$$

is a nontrivial 4-srm polynomial which is not divided by $x^2 - 4$ but $f(2) = 0$ and

$$g(x) = x^6 + 3x^4 + 3x^3 + 2x^2 + 4 \in \mathbf{F}_5[x]$$

is a nontrivial 4-srm polynomial which is not divided by $x^2 - 4$ but $g(3) = 0$.

Theorem 4 *Let $f(x) \in \mathbf{F}_q[x]$ be a nontrivial a-srm polynomial of degree $n = 2m$ and a be a square in $\mathbf{F}_q$. If $f(x)$ is not divided by $x^2 - a$ and $f\left(\sqrt{a}\right) = 0$, then it can be written as $f(x) = \left(x - \sqrt{a}\right)^k \cdot g(x)$, where k is even and $g(x)$ is a nontrivial a-srm polynomial with $g\left(\sqrt{a}\right) \neq 0$.*

Proof. Let $f(x) = \sum_{i=0}^{n} b_i x^i$ then by assumption $b_0 = f(0) = a^m = \sqrt{a}^n$. Set

$$f_1(x) := \frac{f(x)}{x - \sqrt{a}} = x^{n-1} + c_{n-2}x^{n-2} + \cdots + c_1 x + c_0 \in \mathbf{F}_q[x].$$

Clearly $(-\sqrt{a}) \cdot c_0 = b_0 = \sqrt{a}^n$ which implies that $c_0 = -\sqrt{a}^{n-1}$. Then $f_1(x)$ is a-srm polynomial since $x - \sqrt{a}$ is a-srm polynomial. By Theorem 2, $f_1(x)$ is divided by $x - \sqrt{a}$ and thus we can write as $f(x) = \left(x - \sqrt{a}\right)^2 \cdot f_2(x)$. It is clear that $\left(x - \sqrt{a}\right)^2$ is a nontrivial a-srm polynomial, hence $f_2(x)$ is a nontrivial a-srm polynomial of degree $n - 2$. Continuing this procedure for $f_2(x)$ complete the proof. □

When a is a square in $\mathbf{F}_q$, from this theorem the consideration of nontrivial a-srm polynomials not divided by $x^2 - a$ is reduced to the case of ones without roots $\sqrt{a}$ and $-\sqrt{a}$.

3 The number of a-srim polynomials over finite fields

In this section we generalize some results of Section 2.7 in [6]. For a positive integer n, denote $H_{n,q}(x) := x^{q^n+1} - a$.

Theorem 5 *If a is a square in $\mathbf{F}_q$, then $H_{n,q}(x)$ is divided by $x^2 - a$ and if a is not a square in $\mathbf{F}_q$, then $H_{n,q}(x)$ is divided by $x^2 - a$ if and only if n is even.*

Proof. If a is a square in $\mathbf{F}_q$, then $\sqrt{a}, -\sqrt{a} \in \mathbf{F}_q$ and

$$H_{n,q}\left(\sqrt{a}\right) = \sqrt{a}^{q^n+1} - \sqrt{a}^2 = \sqrt{a}\left(\sqrt{a}^{q^n} - \sqrt{a}\right) = 0$$

for $\sqrt{a}^q = \sqrt{a}$ by [7, Lemma 2.2]. Similarly $H_{n,q}\left(-\sqrt{a}\right) = 0$ and thus $H_{n,q}(x)$ is divided by $x^2 - a$. If a is not a square in $\mathbf{F}_q$, then $x^2 - a$ is irreducible over $\mathbf{F}_q$ and by [7, Lemma 2.12, 2.13],

$$x^2 - a | H_{n,q}(x) \Leftrightarrow H_{n,q}\left(\sqrt{a}\right) = 0 \Leftrightarrow \sqrt{a}^{q^n} - \sqrt{a} = 0 \Leftrightarrow 2|n$$

which complete the proof. □

By this theorem, whether $H_{n,q}(x)$ is divided by $x^2 - a$ depends on whether one of the following conditions holds.

a) a is a square in $\mathbf{F}_q$

b) a is not a square in $\mathbf{F}_q$ and n is even.

Theorem 6 (1) *$H_{n,q}(x)$ is divided by any nontrivial a-srim polynomial of degree $2n$ over $\mathbf{F}_q$.*

(2) *Each irreducible factor $f(x)$ of degree ≥ 2 of $H_{n,q}(x)$ such that $f(x) \neq x^2 - a$ is a nontrivial a-srim polynomial of degree $2d$, where $d|n$ and n/d is odd.*

Proof. (1) is clear from [4, Theorem 2.1] and [7, Lemma 2.12]. To prove (2), let α be a root of $f(x)$. Then $\alpha^{q^n+1} = a$, that is, $\alpha^{q^n} = a/\alpha$ which implies that $f(x) = \hat{f_a}(x)$. From Corollary 1, $f(x)$ has even degree $2d$. If $a/\alpha = \alpha$ then $\alpha^2 = a$ and $f(x)$ divides $x^2 - a$, a contradiction. Hence $f(x)$ is a nontrivial a-srim polynomial. Since $f(x)$ is irreducible,

$$\alpha, \alpha^q, \alpha^{q^2}, \cdots, \alpha^{q^{2d-1}}$$

are the different roots of $f(x)$ and thus

$$\alpha^{q^{2n}} = \left(\alpha^{q^n}\right)^{q^n} = \left(\frac{a}{\alpha}\right)^{q^n} = \alpha$$

which implies that $\alpha \in \mathbf{F}_{q^{2n}}$. From the definition of conjugates, $2d|2n$ hence $d|n$. But from (1), $f(x)$ is a divisor of $H_{d,q}(x)$. Now suppose that n/d is even, that is, $n = 2kd$. Then $\alpha^{q^{2d}} = \alpha$ and

$$\begin{aligned} 0 = \alpha^{q^n+1} - a = \alpha \cdot \alpha^{q^{2kd}} - a = \alpha \cdot \left(\alpha^{q^{2d}}\right)^{q^{2(k-1)d}} - a \\ = \alpha \cdot \alpha^{q^{2(k-1)d}} - a = \cdots = \alpha \cdot \alpha^{q^{2d}} - a = \alpha^2 - a, \end{aligned}$$

which would contradict the assumption. Therefore n/d is odd. □

We use the similar notation as in [6] for a-srim polynomials , that is, $SI_{n,q}(x)$ and $si(n,q)$ are respectively the product and the number of all a-srim polynomials of degree $2n$ over $\mathbf{F}_q$ and

$$M_{n,q}(x) = \begin{cases} H_{n,q}(x)/(x^2 - a), a) \text{ or } b) \text{ holds}, \\ H_{n,q}(x), \text{otherwise} \end{cases} \quad \delta = \begin{cases} -1, a) \text{ or } b) \text{ holds}, \\ 1, \text{otherwise} \end{cases}$$

Under these notations, we have the following easily.

Corollary 2

$$M_{n,q}(x) = \prod_{\substack{d|n \\ n/d \text{ odd}}} SI_{n,q}(x), \qquad q^n + \delta = \sum_{\substack{d|n \\ n/d \text{ odd}}} 2d \cdot si(d,q)$$

Now consider a formula for the number of all nontrivial a-srim polynomials of degree $2n$ over $\mathbf{F}_q$.

Lemma 3 [6, *Theorem* 2.7.2] *Let f and F be two functions from* $\mathbf{N}$ *to an additive group G. Then the following conditions are equivalent.*

$$(1)F(n) = \sum_{\substack{d|n\\ n/d \ odd}} f(d) \quad (\forall n \in \mathbf{N}) \quad (2)f(n) = \sum_{\substack{d|n\\ d \ odd}} \mu(d)F(n/d) \quad (\forall n \in \mathbf{N})$$

where μ is Möbius function.

Let G be a multiplicative group of all rational functions over $\mathbf{F}_q$, and set

$$F(n) = M_{n,q}(x), \qquad f(n) = SI_{n,q}(x).$$

Then from above lemma and corollary we have

$$SI_{n,q}(x) = \prod_{\substack{d|n\\ d \text{ odd}}} M_{n/d,q}(x)^{\mu(d)} \tag{2}$$

$$si(n,q) = \frac{1}{2n} \sum_{\substack{d|n\\ d \text{ odd}}} \mu(d) \left(q^{n/d} + \delta\right) \tag{3}$$

Theorem 7 *If $n = 1$ then*

$$si(n,q) = \begin{cases} (q-1)/2, & a \ is \ a \ square \ in \ \mathbf{F}_q \\ (q+1)/2, & otherwise \end{cases} \tag{4}$$

and if $n > 1$ then

$$si(n,q) = \begin{cases} (q^n - 1)/(2n), & n = 2^k \ (k \in \mathbf{N}) \\ \frac{1}{2n} \sum_{\substack{d|n\\ d \ odd}} \mu(d) q^{n/d}, & otherwise \end{cases} \tag{5}$$

Proof. If $n = 1$ then b) can not hold. So $\delta = -1$ if a is a square in $\mathbf{F}_q$ and $\delta = 1$ otherwise, hence (4) follows from (3). On the other hand, it follows

$$\sum_{\substack{d|n\\ d \text{ odd}}} \mu(d) = \begin{cases} 1, & n = 2^k \ (k \text{ is nonnegative integer}) \\ 0, & \text{otherwise} \end{cases} \tag{6}$$

from the properties of Möbius function. If $n > 1$ then we have from (3)

$$si(n,q) = \frac{1}{2n} \sum_{\substack{d|n\\ d \text{ odd}}} \mu(d) q^{n/d} + \frac{\delta}{2n} \sum_{\substack{d|n\\ d \text{ odd}}} \mu(d) \tag{7}$$

If $n = 2^k \ (k \in \mathbf{N})$, then $si(n,q) = (q^n - 1)/2n$ because 1 is unique odd divisor of n and $\delta = -1$. In other cases the claim immediately follows from (6) and (7). $\square$

4 The parity of the number of irreducible factors of generalized self-reciprocal polynomials

Theorem 8 *Let $f(x) \in \mathbf{F}_q[x]$ be a nontrivial a-srm polynomial of degree $2n$ with r pairwise distinct irreducible factors over $\mathbf{F}_q$. Then $r \equiv 0 \pmod 2$ if and only if $(-1)^n a^{n(n-2)} f(\sqrt{a})\, f(-\sqrt{a})$ is a square in $\mathbf{F}_q$, where $\sqrt{a}$ is a square of a in some extension of $\mathbf{F}_q$.*

Proof. Let $f(x) = \sum_{i=0}^{2n} b_i x^i$, then $f(x) = x^n g\left(x + \frac{a}{x}\right)$, where $g(x) = b_n + \sum_{i=0}^{n-1} b_{2n-i} D_{n-i,a}(x)$ and $D_{n-i,a}(x)$ is the Dickson polynomial [4]. Clearly $f(0) = a^n \neq 0$ and by properties of the resultant, $R(f, f') = R(f, nf - xf')$. Since

$$\begin{aligned} f'(x) &= nx^{n-1} g(x + a/x) + x^n g'(x + a/x)(1 - a/x^2) \\ &= nx^{n-1} g(x + a/x) + x^{n-2} g'(x + a/x)(x^2 - a) \end{aligned}$$

and $nf(x) - xf'(x) = -x^{n-1} g'(x + a/x)(x^2 - a)$, we have

$$\begin{aligned} D(f) = (-1)^{n(2n-1)} R(f, f') &= (-1)^n \cdot f(0)^{-1} \cdot R(f, nf - xf') \\ &= (-1)^n \cdot a^{-n} \cdot R(f, x^2 - a) \cdot R(f, -x^{n-1} g'(x + a/x)) \\ &= (-1)^n \cdot a^{-n} \cdot f\left(\sqrt{a}\right) f\left(-\sqrt{a}\right) \cdot R(f, -x^{n-1} g'(x + a/x)) \end{aligned}$$

Let $x_0, \cdots, x_{n-1}, a/x_0, \cdots, a/x_{n-1}$ be the roots of $f(x)$ in some extension of $\mathbf{F}_q$. Then

$$\begin{aligned} R\left(f, -x^{n-1} g'\left(x + \frac{a}{x}\right)\right) &= \prod_{i=0}^{n-1} x_i^{n-1} g'\left(x_i + \frac{a}{x_i}\right) \prod_{i=0}^{n-1} \left(\frac{a}{x_i}\right)^{n-1} g'\left(\frac{a}{x_i} + x_i\right) \\ &= \prod_{i=0}^{n-1} x_i^{n-1} g'\left(x_i + \frac{a}{x_i}\right) \prod_{i=0}^{n-1} a^{n-1} x_i^{1-n} g'\left(x_i + \frac{a}{x_i}\right) \\ &= a^{n(n-1)} \left[\prod_{i=0}^{n-1} g'\left(x_i + \frac{a}{x_i}\right)\right]^2. \end{aligned}$$

On the other hand, since x_i is a root of $f(x)$ if and if $x_i + a/x_i$ is a root of $g(x)$ and

$$D(g) = (-1)^{n(n-1)/2} R(g, g') = (-1)^{n(n-1)/2} \prod_{i=0}^{n-1} g'\left(x_i + \frac{a}{x_i}\right),$$

we have

$$D(f) = (-1)^n a^{n(n-2)} f\left(\sqrt{a}\right) f\left(-\sqrt{a}\right) \cdot D(g)^2.$$

Required result follows from Stickelberger theorem [9] (See also [1, Theorem 4]). $\square$

Let $f(x)=\sum_{i=0}^{2n} b_i x^i$ be as in Theorem 8, then

$$f(x)=\sum_{i=0}^{n} b_{2i}x^{2i}+\sum_{i=0}^{n-1} b_{2i+1}x^{2i+1}$$

and

$$f\left(\sqrt{a}\right)=\sum_{i=0}^{n} b_{2i}a^{i}+\sqrt{a}\sum_{i=0}^{n-1} b_{2i+1}a^{i},$$
$$f\left(-\sqrt{a}\right)=\sum_{i=0}^{n} b_{2i}a^{i}-\sqrt{a}\sum_{i=0}^{n-1} b_{2i+1}a^{i}.$$

Denote

$$A:=\sum_{i=0}^{n} b_{2i}a^{i},\qquad B:=\sum_{i=0}^{n-1} b_{2i+1}a^{i}$$

then $f\left(\sqrt{a}\right)f\left(-\sqrt{a}\right)=A^2-aB^2$, which means $f\left(\sqrt{a}\right)f\left(-\sqrt{a}\right)\in \mathbf{F}_q$. Since $f(x)$ is nontrivial, $b_i=b_{n-i}a^{n-i}$. Therefore if n is odd then

$$A=2\sum_{i=0}^{(n-1)/2} b_{2n-2i}a^{n-i},\quad B=2\sum_{i=0}^{(n-3)/2} b_{2n-2i-1}a^{n-i-1}+b_n a^{(n-1)/2}.$$

and if n is even then

$$A=2\sum_{i=0}^{n/2-1} b_{2n-2i}a^{n-i}+b_n a^{n/2},\quad B=2\sum_{i=0}^{n/2-1} b_{2n-2i-1}a^{n-i-1}.$$

For any monic polynomial $f(x)\in \mathbf{F}_q[x]$ of degree n, denote $f_a^Q(x):=x^n f(x+a/x)$. Then $f_a^Q(x)$ is a nontrivial a-srm polynomial of degree $2n$. This is a natural generalization of $f^Q(x)$ in [8] and it has been mentioned in [4], too.

Theorem 9 *Suppose that $f(x)\in \mathbf{F}_q[x]$ is a monic irreducible polynomial of degree $n(\geq 1)$ and $f\left(\sqrt{a}\right)f\left(-\sqrt{a}\right)\neq 0$. Then $f_a^Q(x)$ is either a-srim polynomial or a product of a-reciprocal pair of irreducible polynomials of degree n which are not a-self reciprocal.*

Proof. Let α be any root of $f_a^Q(x)$. Then $\beta=\alpha+a/\alpha$ is a root of $f(x)$, so $\beta^{q^n}=\beta$. Here n is the smallest positive integer k with $\beta^{q^k}=\beta$. Multiply α^{q^n} to both sides of the identity

$$\alpha^{q^n}+\frac{a}{\alpha^{q^n}}=\beta^{q^n}=\beta=\alpha+\frac{a}{\alpha},$$

then

$$\alpha^{2q^n} + a = \alpha^{q^n+1} + a\alpha^{q^n-1}$$

and so

$$\left(\alpha^{q^n+1} - a\right)\left(\alpha^{q^n-1} - 1\right) = 0.$$

Hence $\alpha^{q^n+1} = a$ or $\alpha^{q^n-1} = 1$. When $\alpha^{q^n+1} = a$, the irreducible factor $g(x)$ of $f_a^Q(x)$ with a root α had degree ≥ 2 because if $g(x)$ has degree 1, then a is a square in $\mathbf{F}_q$ and $g(x) = x \pm \sqrt{a}$ which contradict to $f\left(\sqrt{a}\right) f\left(-\sqrt{a}\right) \neq 0$. Therefore $g(x)$ is a nontrivial a-srm polynomial of degree $2d$ for a positive divisor of n. If we assume $d < n$ then $\alpha^{q^d+1} = a$, so $\beta^{q^d} = \beta$, a contradiction. Hence $d = n$ and $f_a^Q(x)$ is irreducible over $\mathbf{F}_q$. When $\alpha^{q^n-1} = 1$, the irreducible factor $g(x)$ of $f_a^Q(x)$ with a root α divides $x^{q^n} - x$ and the degree of $g(x)$ is n. Therefore $f_a^Q(x)$ is a product of two n-degree irreducible polynomials $g(x)$ and $h(x) = f_a^Q(x)/g(x)$. Suppose $g(x)$ is a nontrivial a-srim polynomial, then $\alpha^{q^{n/2}+1} - a = 0$ which contradicts to the minimality of n . So $g(x)$ and $h(x)$ are not a-self reciprocal. And if $f_a^Q(x)$ has a root α then it also has a root a/α, which implies easily that $h(x)$ is a-reciprocal of $g(x)$. □

Finally we consider whether Theorem 8 is valid if $f(x)$ has repeated irreducible factors.

Theorem 10 *Let $f(x) \in \mathbf{F}_q[x]$ be a nontrivial a-srm polynomial of degree $2n$ with $f\left(\sqrt{a}\right) f\left(-\sqrt{a}\right) \neq 0$, and let r be the number of irreducible factors counted with multiplicity of $f(x)$. Then r is even if and only if $(-1)^n a^{n(n-2)} f\left(\sqrt{a}\right) f\left(-\sqrt{a}\right)$ is a square in $\mathbf{F}_q$.*

Proof. Like as in the proof of Theorem 8, there exists $g(x) \in \mathbf{F}_q[x]$ such that $f(x) = x^n g(x + a/x)$. Suppose that $g(x) = g_1(x) \cdots g_k(x)$ where $g_i(x) \in \mathbf{F}_q[x]$ is monic irreducible polynomial of degree n_i and $n_1 + \cdots + n_k = n$. Denote $f_i(x) = x^{n_i} g(x + a/x)$, then $f(x) = f_1(x) \cdots f_k(x)$ where every $f_i(x)$ is a nontrivial a-srm polynomial of degree $2n_i$ over $\mathbf{F}_q$. By Theorem 9, every $f_i(x)$ is either irreducible or a product of two distinct monic irreducible polynomials of degree n_i. Now suppose that the claim holds for two nontrivial a-srm polynomials $g(x)$, $h(x)$ of degree $2s$, $2t$ with r_s, r_h irreducible factors counted with multiplicity respectively over $\mathbf{F}_q$. If both r_s and r_h are odd, then neither $(-1)^s a^{s(s-2)} g\left(\sqrt{a}\right) g\left(-\sqrt{a}\right)$ nor $(-1)^t a^{t(t-2)} h\left(\sqrt{a}\right) h\left(-\sqrt{a}\right)$ is a square in $\mathbf{F}_q$, hence $(-1)^{s+t} a^{(s+t)(s+t-2)} g\left(\sqrt{a}\right) g\left(-\sqrt{a}\right) h\left(\sqrt{a}\right) h\left(-\sqrt{a}\right)$ is a square in $\mathbf{F}_q$. Similar observation about other cases for r_s and r_h tell us the claim is also valid for $g(x)h(x)$. It is sufficient to apply Theorem 8 to complete the proof. □

References

[1] O.Ahmadi, Self-reciprocal irreducible pentanomials over $\mathbf{F}_2$. Des. Codes Cryptogr, 38(2006) 395-397

[2] O.Ahmadi G.Vega, On the parity of the number of irreducible factors of self-reciprocal polynomials over finite fields, Finite Fields Appl. 14(2008) 124-131

[3] L.Carlitz, Some theorems on irreducible reciprocal polynomials over a finite field, J. reine Angew. Math. 227(1967) 212-220

[4] R.W.Fitzgerald J.L.Yucas, Generalized reciprocals, factors of Dickson polynomials and generalized cyclotomic polynomials over finite fields, Finite Fields Appl. 13(2007) 492-515

[5] T.Garefalakis, Self-reciprocal irreducible polynomials with prescribed coefficients, Finite Fields Appl. 17(2011) 183-193

[6] D.Jungnickel, Finite Fields, Structure and Arithmetic, BI-Wissenschaftverlag, Manheim, 1993

[7] R.Lidl H.Niederreiter, Finite Fields, Cambridge Univ. Press, Cambridge, 1997

[8] H.Meyn, On the construction of irreducible self-reciprocal polynomials over finite fields, Appl. Algebra Engrg. Comm. Comput. 1(1990), 43-53

[9] L.Stickelberger, Über eine neue Eigenschaft der Diskriminanten algebraischer Zahlk örper, Verhandlungen des ersten Internationalen Mathematiker-Kongresses, Zürich,(1897) 182-193

[10] J.L.Yucas G.L.Mullen, Self-reciprocal irreducible polynomials over finite fields, Des. Codes Cryptogr. 33(2004) 275-281